

Properties of Fuzzy Relation

Noboru Endou
Gifu National College of Technology

Takashi Mitsuishi
Miyagi University

Keiji Ohkubo
Shinshu University
Nagano

Summary. In this article, we introduce four fuzzy relations and the composition, and some useful properties are shown by them. In section 2, the definition of converse relation R^{-1} of fuzzy relation R and properties concerning it are described. In the next section, we define the composition of the fuzzy relation and show some properties. In the final section we describe the identity relation, the universe relation and the zero relation.

MML Identifier: FUZZY_4.

The notation and terminology used here are introduced in the following papers: [5], [6], [2], [9], [4], [3], [8], [7], and [1].

1. BASIC PROPERTIES OF THE MEMBERSHIP FUNCTION

We follow the rules: x, y, z are sets and C_1, C_2, C_3 are non empty sets.

Let C_1 be a non empty set and let F be a membership function of C_1 . One can check that $\text{rng } F$ is non empty.

Next we state four propositions:

- (1) Let F be a membership function of C_1 . Then $\text{rng } F$ is bounded and for every x such that $x \in \text{dom } F$ holds $F(x) \leq \sup \text{rng } F$ and for every x such that $x \in \text{dom } F$ holds $F(x) \geq \inf \text{rng } F$.
- (2) For all membership functions F, G of C_1 such that for every x such that $x \in C_1$ holds $F(x) \leq G(x)$ holds $\sup \text{rng } F \leq \sup \text{rng } G$.

- (3) For every Membership function f of C_1, C_2 and for every element c of $[C_1, C_2]$ holds $0 \leq f(c)$ and $f(c) \leq 1$.
- (4) For every Membership function f of C_1, C_2 and for all x, y such that $\langle x, y \rangle \in [C_1, C_2]$ holds $0 \leq f(\langle x, y \rangle)$ and $f(\langle x, y \rangle) \leq 1$.

2. DEFINITION OF CONVERSE FUZZY RELATION AND SOME PROPERTIES

Let C_1, C_2 be non empty sets and let h be a Membership function of C_2, C_1 . The functor converse h yielding a Membership function of C_1, C_2 is defined by:

(Def. 1) For all x, y such that $\langle x, y \rangle \in [C_1, C_2]$ holds $(\text{converse } h)(\langle x, y \rangle) = h(\langle y, x \rangle)$.

Let C_1, C_2 be non empty sets, let f be a Membership function of C_2, C_1 , and let R be a fuzzy relation of C_2, C_1, f . The functor R^{-1} yields a fuzzy relation of C_1, C_2 , converse f and is defined by:

(Def. 2) $R^{-1} = [[C_1, C_2], (\text{converse } f)^\circ [C_1, C_2]]$.

The following propositions are true:

- (5) For every Membership function f of C_1, C_2 holds $\text{converse } \text{converse } f = f$.
- (6) For every Membership function f of C_1, C_2 and for every fuzzy relation R of C_1, C_2, f holds $(R^{-1})^{-1} = R$.
- (7) For every Membership function f of C_1, C_2 holds $1 - \text{converse } f = \text{converse } (1 - f)$.
- (8) For every Membership function f of C_1, C_2 and for every fuzzy relation R of C_1, C_2, f holds $(R^{-1})^c = (R^c)^{-1}$.
- (9) For all Membership functions f, g of C_1, C_2 holds $\text{converse } \max(f, g) = \max(\text{converse } f, \text{converse } g)$.
- (10) Let f, g be Membership functions of C_1, C_2, R be a fuzzy relation of C_1, C_2, f , and S be a fuzzy relation of C_1, C_2, g . Then $(R \cup S)^{-1} = R^{-1} \cup S^{-1}$.
- (11) For all Membership functions f, g of C_1, C_2 holds $\text{converse } \min(f, g) = \min(\text{converse } f, \text{converse } g)$.
- (12) Let f, g be Membership functions of C_1, C_2, R be a fuzzy relation of C_1, C_2, f , and S be a fuzzy relation of C_1, C_2, g . Then $(R \cap S)^{-1} = R^{-1} \cap S^{-1}$.
- (13) Let f, g be Membership functions of C_1, C_2 and given x, y . If $x \in C_1$ and $y \in C_2$, then if $f(\langle x, y \rangle) \leq g(\langle x, y \rangle)$, then $(\text{converse } f)(\langle y, x \rangle) \leq (\text{converse } g)(\langle y, x \rangle)$.
- (14) Let f, g be Membership functions of C_1, C_2, R be a fuzzy relation of C_1, C_2, f , and S be a fuzzy relation of C_1, C_2, g . If $R \subseteq S$, then $R^{-1} \subseteq S^{-1}$.

- (15) For all Membership functions f, g of C_1, C_2 holds
 $\text{converse } \min(f, 1-\text{minus } g) = \min(\text{converse } f, 1-\text{minus } \text{converse } g)$.
- (16) Let f, g be Membership functions of C_1, C_2, R be a fuzzy relation of C_1, C_2, f , and S be a fuzzy relation of C_1, C_2, g . Then $(R \setminus S)^{-1} = R^{-1} \setminus S^{-1}$.
- (17) For all Membership functions f, g of C_1, C_2 holds converse
 $\max(\min(f, 1-\text{minus } g), \min(1-\text{minus } f, g)) =$
 $\max(\min(\text{converse } f, 1-\text{minus } \text{converse } g),$
 $\min(1-\text{minus } \text{converse } f, \text{converse } g))$.
- (18) Let f, g be Membership functions of C_1, C_2, R be a fuzzy relation of C_1, C_2, f , and S be a fuzzy relation of C_1, C_2, g . Then $(R \dot{\setminus} S)^{-1} = R^{-1} \dot{\setminus} S^{-1}$.

3. DEFINITION OF THE COMPOSITION AND SOME PROPERTIES

Let C_1, C_2, C_3 be non empty sets, let h be a Membership function of C_1, C_2 , let g be a Membership function of C_2, C_3 , and let x, z be sets. Let us assume that $x \in C_1$ and $z \in C_3$. The functor $\min(h, g, x, z)$ yields a membership function of C_2 and is defined by:

- (Def. 3) For every element y of C_2 holds $(\min(h, g, x, z))(y) = \min(h(\langle x, y \rangle), g(\langle y, z \rangle))$.

Let C_1, C_2, C_3 be non empty sets, let h be a Membership function of C_1, C_2 , and let g be a Membership function of C_2, C_3 . The functor hg yields a Membership function of C_1, C_3 and is defined by:

- (Def. 4) For all x, z such that $\langle x, z \rangle \in [C_1, C_3]$ holds $(hg)(\langle x, z \rangle) = \sup \text{rng } \min(h, g, x, z)$.

Let C_1, C_2, C_3 be non empty sets, let f be a Membership function of C_1, C_2 , let g be a Membership function of C_2, C_3 , let R be a fuzzy relation of C_1, C_2, f , and let S be a fuzzy relation of C_2, C_3, g . The functor RS yields a fuzzy relation of C_1, C_3, fg and is defined as follows:

- (Def. 5) $RS = [[C_1, C_3], (fg)^\circ [C_1, C_3]]$.

Next we state a number of propositions:

- (19) For every Membership function f of C_1, C_2 and for all Membership functions g, h of C_2, C_3 holds $f \max(g, h) = \max(fg, fh)$.
- (20) Let f be a Membership function of C_1, C_2, g, h be Membership functions of C_2, C_3, R be a fuzzy relation of C_1, C_2, f, S be a fuzzy relation of C_2, C_3, g , and T be a fuzzy relation of C_2, C_3, h . Then $R(S \cup T) = RS \cup RT$.
- (21) For all Membership functions f, g of C_1, C_2 and for every Membership function h of C_2, C_3 holds $\max(f, g)h = \max(fh, gh)$.

- (22) Let f, g be Membership functions of C_1, C_2 , h be a Membership function of C_2, C_3 , R be a fuzzy relation of C_1, C_2 , f, S be a fuzzy relation of C_1, C_2, g , and T be a fuzzy relation of C_2, C_3, h . Then $(R \cup S)T = RT \cup ST$.
- (23) Let f be a Membership function of C_1, C_2 , g, h be Membership functions of C_2, C_3 , and x, z be sets. If $x \in C_1$ and $z \in C_3$, then $(f \min(g, h))(\langle x, z \rangle) \leq (\min(fg, fh))(\langle x, z \rangle)$.
- (24) Let f be a Membership function of C_1, C_2 , g, h be Membership functions of C_2, C_3 , R be a fuzzy relation of C_1, C_2 , f, S be a fuzzy relation of C_2, C_3 , g , and T be a fuzzy relation of C_2, C_3, h . Then $R(S \cap T) \subseteq (RS) \cap (RT)$.
- (25) Let f, g be Membership functions of C_1, C_2 , h be a Membership function of C_2, C_3 , and x, z be sets. If $x \in C_1$ and $z \in C_3$, then $(\min(f, g)h)(\langle x, z \rangle) \leq (\min(fh, gh))(\langle x, z \rangle)$.
- (26) Let f, g be Membership functions of C_1, C_2 , h be a Membership function of C_2, C_3 , R be a fuzzy relation of C_1, C_2 , f, S be a fuzzy relation of C_1, C_2 , g , and T be a fuzzy relation of C_2, C_3, h . Then $(R \cap S)T \subseteq (RT) \cap (ST)$.
- (27) For every Membership function f of C_1, C_2 and for every Membership function g of C_2, C_3 holds converse $fg =$ converse g converse f .
- (28) Let f be a Membership function of C_1, C_2 , g be a Membership function of C_2, C_3 , R be a fuzzy relation of C_1, C_2 , f , and S be a fuzzy relation of C_2, C_3, g . Then $(RS)^{-1} = S^{-1}R^{-1}$.
- (29) Let f, g be Membership functions of C_1, C_2 , h, k be Membership functions of C_2, C_3 , and x, z be sets. Suppose $x \in C_1$ and $z \in C_3$ and for every set y such that $y \in C_2$ holds $f(\langle x, y \rangle) \leq g(\langle x, y \rangle)$ and $h(\langle y, z \rangle) \leq k(\langle y, z \rangle)$. Then $(fh)(\langle x, z \rangle) \leq (gk)(\langle x, z \rangle)$.
- (30) Let f, g be Membership functions of C_1, C_2 , h, k be Membership functions of C_2, C_3 , R be a fuzzy relation of C_1, C_2 , f, S be a fuzzy relation of C_1, C_2, g , T be a fuzzy relation of C_2, C_3, h , and W be a fuzzy relation of C_2, C_3, k . If $R \subseteq S$ and $T \subseteq W$, then $RT \subseteq SW$.

4. DEFINITION OF IDENTITY RELATION AND PROPERTIES OF UNIVERSE AND ZERO RELATION

Let C_1, C_2 be non empty sets. The functor $\text{Imf}(C_1, C_2)$ yields a Membership function of C_1, C_2 and is defined as follows:

- (Def. 6) For all x, y such that $\langle x, y \rangle \in [C_1, C_2]$ holds if $x = y$, then $(\text{Imf}(C_1, C_2))(\langle x, y \rangle) = 1$ and if $x \neq y$, then $(\text{Imf}(C_1, C_2))(\langle x, y \rangle) = 0$.

One can prove the following propositions:

- (31) For every element c of $[C_1, C_2]$ holds $(\text{Zmf}(C_1, C_2))(c) = 0$ and $(\text{Umf}(C_1, C_2))(c) = 1$.

- (32) For all x, y such that $\langle x, y \rangle \in [C_1, C_2]$ holds $(Zmf(C_1, C_2))(\langle x, y \rangle) = 0$ and $(Umf(C_1, C_2))(\langle x, y \rangle) = 1$.
- (33) Let f be a Membership function of C_2, C_3 , O_1 be a zero relation of C_1, C_2 , O_2 be a zero relation of C_1, C_3 , and R be a fuzzy relation of C_2, C_3, f . Then $O_1 R = O_2$.
- (34) For every Membership function f of C_1, C_2 holds $f Zmf(C_2, C_3) = Zmf(C_1, C_3)$.
- (35) Let f be a Membership function of C_1, C_2 , O_1 be a zero relation of C_2, C_3 , O_2 be a zero relation of C_1, C_3 , and R be a fuzzy relation of C_1, C_2, f . Then $R O_1 = O_2$.
- (36) For every Membership function f of C_1, C_1 holds $f Zmf(C_1, C_1) = Zmf(C_1, C_1) f$.
- (37) Let f be a Membership function of C_1, C_1 , O be a zero relation of C_1, C_1 , and R be a fuzzy relation of C_1, C_1, f . Then $R O = O R$.

REFERENCES

- [1] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [2] Czesław Byliński and Piotr Rudnicki. Bounding boxes for compact sets in \mathcal{E}^2 . *Formalized Mathematics*, 6(3):427–440, 1997.
- [3] Jarosław Kotowicz. Convergent real sequences. Upper and lower bound of sets of real numbers. *Formalized Mathematics*, 1(3):477–481, 1990.
- [4] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [5] Takashi Mitsuishi, Noboru Endou, and Yasunari Shidama. The concept of fuzzy set and membership function and basic properties of fuzzy set operation. *Formalized Mathematics*, 9(2):351–356, 2001.
- [6] Takashi Mitsuishi, Katsumi Wasaki, and Yasunari Shidama. The concept of fuzzy relation and basic properties of its operation. *Formalized Mathematics*, 9(3):517–524, 2001.
- [7] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [8] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [9] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.

Received June 25, 2001
